

US and Scottish teens reflections on sexual risk reduction: Common healthcare intervention challenges across continents

Dianne Morrison-Beedy, PhD, RN, WHNP, FNAP, FAANP, FAAN
Chief Talent & Global Strategy Officer and
The Centennial Professor of Nursing
The Ohio State University
USA



THE OHIO STATE UNIVERSITY

COLLEGE OF NURSING

It is hoped that the information provided in this overview will be helpful to attendees in familiarizing themselves with the research topic. This overview is not intended to be comprehensive nor does it involve sales of a product. Subjects covered in this presentation represent my views and not those of the any university.



Acknowledgements

- NIH/NINR Maintaining HIV Prevention Gains in Female Adolescents RO1NR008194
- NIH/NINR Maintaining abstinence and reducing HIV in adolescent girls R03NR010193
- NIH/NINR Motivational HIV prevention intervention for young women KO1NR000152
- NIH/NINR HIV risk appraisal in single urban women R15NR04059
- NIH/NIMH Community adults and parents response to HIV prevention intervention for young adolescents P30A1078498
- Also University of Rochester, Syracuse University, Niagara University, Sigma Theta Tau International, Helping People with AIDS, AWHONN and others
- And to my team and co-Is Drs. Michael P Carey, Xin Tu, project director Sheryl Jones and the best team ever



Background


Impact of sexual risk behaviors in adolescent girls across the globe

- 16 million girls aged 15-19 and 1 million girls < 15 give birth every year
- Yearly >300 million new cases of curable STIs occur worldwide with the highest rates in 15-24 years old
- Globally, > 90% of HIV is transmitted heterosexually, majority in females
- Impacts: high school dropout, welfare, healthcare costs, lower wages



Adolescent Females: The Disproportionate Impact of HIV

Percentage of AIDS cases world wide





Components of Successful Interventions

Theoretically-driven

Tailored to specific population, culturally relevant

Focus beyond knowledge to skill enhancement and motivation

Intervention implemented in accessible points in communities

Optimizing uptake :Reproducibility, accessibility



HIPTeens Step by Step


Step 1. Identify population's existing level of HIV-prevention IMB

Step 2. Design population-specific intervention strategies

Step 3. Methodologically rigorous evaluation research




HIPTeens RCT



* STI Test




























National Recognition of *HIPTeens*

CDC

- Has been identified by the CDC as meeting PRS criteria for good evidence of efficacy HIV & STI prevetion
 - disseminated through the CDC
 - http://www.cdc.gov/hiv/prevention/researc
 h/compendium/rr/complete.html

HHS

- Recognized by U.S. Health and Human Services as meeting intervention effectiveness criteria based on the Teen Pregnancy Prevention (TPP) Evidence Review
- www.hhs.gov/ash/oah/oahinitiatives/teen_pregnancy/db/programs/hip_t eens.pdf



Purpose: To analyze an integrated data set from multiple studies conducted in the formative work for this intervention with sexually-active and abstinent teens of both genders in two countries.

Goal: To identify common themes and threads addressing risk-taking and protective behaviors.



Participants and Recruitment

- Three studies: 72 adolescents ages 14-19 years
- Enrolled both sexually active and abstinent participants of both genders the majority of whom were economically-disadvantaged young people of color (64%)
- Majority female (85%) and sexually-active (58%)
- Recruited from after-school programs, community teen-based clinics and social agencies, an urban community "cafe" and by word-of-mouth

THE OHIO STATE UNIVERSITY

Methods: Formative research using focus groups

Followed standard methodological protocols:

- trained moderator
- focus group script
- sessions taped and transcribed verbatim

Analysis: We conducted within- and betweengroup content analysis, identifying and confirming predominant codes and themes.



Knowing the facts is different than knowing how







COLLEGE OF NURSING

Communication and negotiation skills are critical gaps

Let's talk about sex, baby
Let's talk about you and me
Let's talk about all the good things
And the bad things that may be
Let's talk about sex
Let's talk about sex

Salt-N-Pepa







Risky sex often occurs in public or unfamiliar places and with concurrent use of substances









Teens have significant stress over remaining abstinent or practicing safe

sex.





Findings specific to girls

There is always some risk to self and health with sex

In discussions regarding the use of gender-specific interventions, girls clearly indicated they wanted all-girl interventions so they could feel safe and heard





Findings specific to boys

You're always willing to take the chance if you can get the sex

Males said "boys will be boys" in all-male groups but would interact more positively in mixed-gender groups.





Conclusions

Findings provide evidence of continued risk behaviors in teens

- Unique social-behavioral challenges
- Must be integrated into intervention strategy modification



Conclusions

- A continued focus on communication and negotiation skill-building is important for sexual risk reduction interventions.
- Concurrent substance use remains a critical influence on sexual risk taking behaviors and intervention strategies must integrate both behavioral challenges.



Future Work

Understanding the impact of stress, anxiety and depression in teens on evidence-based intervention outcomes, particularly as they impact sexual risk-taking behaviors, is a gap that remains in both intervention and dissemination/implementation science.



Future Work

Consider whether gender-specific interventions benefit one gender more than the other.

Determine best ways to tailor and refine sexual risk reduction interventions for teens



Thank You

Dianne Morrison-Beedy, PhD, RN, WHNP, FNAP, FAANP, FAAN

Chief Talent & Global Strategy Officer

Centennial Professor of Nursing

Fulbright Specialist; Fulbright Scholar UK 2016-2017, Fulbright Scholar Norway 2019

The Ohio State University College of Nursing

Room 223, 760 Kinnear, Columbus, OH 43212

Office 614-514-5808

morrison-beedy.1@osu.edu osu.edu

