

References:

Centers for Disease Control and Prevention (2013). CDC vital signs: Preventing repeat teen births. Retrieved from <http://www.cdc.gov/vitalsigns/teenpregnancy/>

Centers for Disease Control and Prevention (2015). STDs in racial and ethnic minorities. Retrieved from <https://www.cdc.gov/std/stats14/minorities.htm>

Centers for Disease Control and Prevention (2016). HIV among youth. Retrieved from <http://www.cdc.gov/hiv/group/age/youth/index.html>

Cooper, M. L., Shapiro, C. M., & Powers, A. M. (1998). Motivations for sex and risky sexual behavior among adolescents and young adults: a functional perspective. *J Pers Soc Psychol*, 75(6), 1528-1558.

Fisher, J. D., & Fisher, W. A. (1992). Changing AIDS-risk behavior. *Psychological Bulletin*, 111(3), 455-474. doi:10.1037/0033-2909.111.3.455

Gebhardt, W. A., Kuyper, L., & Dusseldorp, E. (2006). Condom use at first intercourse with a new partner in female adolescents and young adults: The Role of cognitive planning and motives for having sex. *Archives of Sexual Behavior*, 35(2), 213.

Gebhardt, W. A., Kuyper, L., & Greunsven, G. (2003). Need for intimacy in relationships and motives for sex as determinants of adolescent condom use. *Journal of Adolescent Health*, 33(3), 154-164. doi:[http://dx.doi.org/10.1016/S1054-139X\(03\)00137-X](http://dx.doi.org/10.1016/S1054-139X(03)00137-X)

Hoffman, S., & Maynard, R. (Eds.). (2008). *Kids having kids: Economic costs and social consequences of teen pregnancy* (2nd ed.). Washington, D.C.: Urban Institute Press.

Jackson, J. M., Seth, P., DiClemente, R. J., & Lin, A. (2015). Association of depressive symptoms and substance use with risky sexual behavior and sexually transmitted infections among african american female adolescents seeking sexual health care. *American Journal of Public Health*, 105(10), 2137-2142. doi:10.2105/AJPH.2014.302493

Kann, L., McManus, T., Harris, W. A., Shanklin, S. L., Flint, K. H., Hawkins, J., & al., e. (2015). Youth risk behavior surveillance — United states, 2015. Atlanta, GA: Centers for Disease Control and Prevention Retrieved from http://www.cdc.gov/healthyyouth/data/yrbs/pdf/2015/ss6506_updated.pdf.

Mazzaferro, K. E., Murray, P. J., Ness, R. B., Bass, D. C., Tyus, N., & Cook, R. L. (2006). Depression, stress, and social support as predictors of high-risk sexual behaviors and STIs in young women. *Journal of Adolescent Health*, 39(4), 601-603. doi:<http://dx.doi.org/10.1016/j.jadohealth.2006.02.004>

Meade, C. S., & Ickovics, J. R. (2005). Systematic review of sexual risk among pregnant and mothering teens in the USA: Pregnancy as an opportunity for integrated prevention of STD and repeat pregnancy. *Social Science & Medicine*, 60(4), 661-678. doi:<http://dx.doi.org/10.1016/j.socscimed.2004.06.015>

Morrison-Beedy, D., Carey, M. P., & Tu, X. (2006). Accuracy of audio computer-assisted self-interviewing (ACASI) and self-administered questionnaires for the assessment of sexual behavior. *AIDS & Behavior*, 10(5), 541.

Morrison-Beedy, D., Jones, S. H., Xia, Y., Tu, X., Crean, H. F., & Carey, M. P. (2013). Reducing sexual risk behavior in adolescent girls: Results from a randomized controlled trial. *Journal of Adolescent Health*, 52(3), 314-321. doi:10.1016/j.jadohealth.2012.07.005

Morrison-Beedy, D., Xia, Y., & Passmore, D. (2013). Sexual risk factors for partner age discordance in adolescent girls and their male partners. *Journal of Clinical Nursing*, 22(23/24), 3289.

Paradise, J. E., Cote, J., Minsky, S., Lourenco, A., & Howland, J. (2001). Personal values and sexual decision-making among virginal and sexually experienced urban adolescent girls. *Journal of Adolescent Health*, 28(5), 404-409. doi:[http://dx.doi.org/10.1016/S1054-139X\(01\)00199-9](http://dx.doi.org/10.1016/S1054-139X(01)00199-9)

Substance Abuse and Mental Health Services Administration. (2012). National survey on drug use and health data spotlight: Depression triples between the ages of 12 and 15 among adolescent girls. Retrieved from <http://www.samhsa.gov/data/sites/default/files/NSDUH-SP77-AdolescentGirlsDepression-2012/CBHSQ-NSDUH-Spotlight-077-AdolescentGirlsDepression-2012.pdf>.

Seth, P., Wingood, G. M., DiClemente, R. J., & Robinson, L. S. (2011). Alcohol use as a marker for risky sexual behaviors and biologically-confirmed sexually transmitted infections among young adult African American women. *Women's health issues*, 21(2), 130-135. doi:10.1016/j.whi.2010.10.005

Shrier, L. A., Walls, C., Lops, C., Kendall, A. D., & Blood, E. A. (2012). Substance use, sexual intercourse, and condom nonuse among depressed adolescents and young adults. *Journal of Adolescent Health*, 50(3), 264-270. doi:<http://dx.doi.org/10.1016/j.jadohealth.2011.07.005>

Vasilenko, S. A., Kugler, K. C., & Rice, C. E. (2016). Timing of first sexual intercourse and young adult health outcomes. *Journal of Adolescent Health*. doi:<http://dx.doi.org/10.1016/j.jadohealth.2016.04.019>

Volpe, E. M., Hardie, T. L., Cerulli, C., Sommers, M. S., & Morrison-Beedy, D. (2013). What's age got to do with it? Partner age difference, power, intimate partner violence, and sexual risk in urban adolescents. *Journal of Interpersonal Violence*, 28(10), 2068-2087. doi:10.1177/0886260512471082