

**References:**

AlGhurair, S. A., Hughes, C. A., Simpson, S. H., & Guirguis, L. M. (2012). A systematic review of patient Self-Reported barriers of adherence to antihypertensive medications using the world health organization multidimensional adherence model. *The Journal of Clinical Hypertension*, 14(12), 877-886.

Al-Maqbali, A. A., Temple-Smith, M., Ferler, J., & Blackberry, I. (2013). Prevalence and determinants of pre-hypertension among omani adults attending non-communicable disease screening program in primary care setting in sohar city. *Oman Medical Journal*, 28(5), 316-323.

Al-Mawali, A. (2015). Non-communicable diseases: Shining a light on cardiovascular disease, oman's biggest killer. *Oman Medical Journal*, 30(4), 227.

Kamran, A., Ahari, S. S., Biria, M., Malpour, A., & Heydari, H. (2015). Determinants of patient's adherence to hypertension medications: Application of health belief model among rural patients. *Annals of Medical and Health Sciences Research*, 4(6), 922-927.

World Health Organization. (2013). A global brief on hypertension, silent killer, global public health crisis. *Retrieve from*

[Http://Apps.Who.Int/Iris/Bitstream/10665/79059/1/WHO\\_DCO\\_WHD\\_2013.2\\_eng.Pdf](Http://Apps.Who.Int/Iris/Bitstream/10665/79059/1/WHO_DCO_WHD_2013.2_eng.Pdf) [Accessed 2013-07-17][WebCite Cache ID 6IBqAI2eN].