

References:

- Boltz, M., Resnick, B., Capezuti, E., Shuluk, J., & Secic, M. (2012). Functional Decline in Hospitalized Older Adults: Can Nursing Make a Difference? *Geriatric Nursing*, 33(4), 272-279. doi:10.1016/j.gerinurse.2012.01.008
- Cruz, A. A., Bousquet, J., & Khaltaev, N. (2007). *Global surveillance, prevention and control of chronic respiratory diseases: a comprehensive approach*: World Health Organization.
- Eaton, T., Kolbe, J., Young, P., Fergusson, W., Moodie, L., Zeng, I., . . . Poole, P. (2009). Does early pulmonary rehabilitation reduce acute health-care utilization in COPD patients admitted with an exacerbation? A randomized controlled study. *Respirology*, 14(2), 230-238. doi:10.1111/j.1440-1843.2008.01418.x
- Greening, N. J., Singh, S. J., Steiner, M. C., Williams, J. E. A., Hussain, S. F., Harvey-Dunstan, T. C., . . . Morgan, M. D. (2014). An early rehabilitation intervention to enhance recovery during hospital admission for an exacerbation of chronic respiratory disease: randomised controlled trial. *British Medical Journal*, 349(5), g4315. doi:10.1136/bmj.g4315
- Healthcare Cost and Utilization Project. (2011). Retrieved from
<https://www.hcup-us.ahrq.gov/reports/statbriefs/sb106.jsp>
- Healthcare Cost and Utilization Project (HCUP). (2011). Retrieved from <http://hcupnet.ahrq.gov/HCUPnet.jsp?Id=FEA5924B0FA108C1&Form=DispTab&GoTo=MAINSEL&JS=Y>
- Liao, L. Y., Chen, K. M., Chung, W. S., & Chien, J. Y. (2015). Efficacy of a respiratory rehabilitation exercise training package in hospitalized elderly patients with acute exacerbation of COPD: a randomized control trial. *International journal of chronic obstructive pulmonary disease*, 10(1), 1703-1709. doi:10.2147/COPD.S90673
- Mundy, L. M., Leet, T. L., Darst, K., Schnitzler, M. A., & Dunagan, W. C. (2003). Early mobilization of patients hospitalized with community-acquired pneumonia. *CHEST Journal*, 124(3), 883-889.
- Nguyen, H. Q., Rondinelli, J., Harrington, A., Desai, S., Liu, I. L. A., Lee, J. S., & Gould, M. K. (2015). Functional status at discharge and 30-day readmission risk in COPD. *Respiratory medicine*, 109(2), 238-246. doi:10.1016/j.rmed.2014.12.004

Nurse-Administered Pulmonary Protocol Increases Out-of-Bed Activity, Shortens Length of Stay, and Reduces Readmissions. (2009). Retrieved from <http://www.innovations.ahrq.gov/content.aspx?id=2425>

Reid, W. D., Camp, P. G., Yamabayashi, C., Goodridge, D., Chung, F., Hunt, M. A., . . . Hoens, A. M. (2012). Exercise prescription for hospitalized people with chronic obstructive pulmonary disease and comorbidities: a synthesis of systematic reviews. *International journal of chronic obstructive pulmonary disease*, 7, 297-320. doi:10.2147/COPD.S29750

Spruit, M. A., Man, W. D. C., Pitta, F., Sewell, L., Raskin, J., Bourbeau, J., . . . Rehabilitation, A. E. T. F. o. P. (2013). An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine*, 188(8), e13-E64. doi:10.1164/rccm.201309-1634ST