

**References:**

- Cardosa, J.M., Kusahara D.M., Guinsburg, R., & Pedreira M.L. (2015). Randomized crossover trial of endotracheal tube suctioning systems in newborn. *Nurs Crit Care*, 40, 1-8.
- Casey H., Newberry, D., Jnah, A. (2016). Early bubble continuous positive airway pressure: investigating interprofessional best practices for the NICU team. *Adv Neon Netw*, 35(3), 125-133.
- Cone, S. (2013). Endotracheal suctioning in preterm infants using four-handed versus routine care. *J Obstet Gynecol Neonatal Nurs*, 42(1), 92-104.
- Mann, B., Sweet, M., Knupp, A., Buck, J., & Chipps, E. (2013). Nasal continuous positive airway pressure: a multisite study of suctioning practices within NICUs. *Adv Neonatal Care* 13(2), E1-9.
- Moretti, C., Gizzi, C., Montecchia, F., Barbara, C., Midulla, F., Sanchez-Luna, M., & Papoff, P. ( 2016). Synchronized nasal intermittent positive pressure ventilation in the newborn and clinical results. *Neonatology* 109(4), 359-65.
- Taheri, P., Asgari, N., Mohammadizadeh, M., & Golchin, M. (2012). The effect of open and closed endotracheal tube suctioning on respiratory parameters of infants undergoing mechanical ventilation. *Iran J Nurse Midwifery Res* 17(10), 26-9.