

References:

Artz, N., Elvers, K. T., Lowe, C. M., Sackley, C., Jepson, P., & Beswick, A. D. (2015). Effectiveness of physiotherapy exercise following total knee

replacement: systematic review and meta-analysis. *BMC Musculoskeletal Disorders*, 16(1), 1.

Collins, N. J., Misra, D., Felson, D. T., Crossley, K. M., & Roos, E. M. (2011). *Measures of knee function: International Knee Documentation Committee*

(IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS),
Knee Injury and Osteoarthritis Outcome Score

Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale
(KOS-ADL), Lysholm Knee Scoring Scale, Oxford

Knee Score (OKS), Western Ontario and McMaster. *Arthritis Care and Research*, 63(SUPPL. 11).
DOI: [10.1002/acr.20632](https://doi.org/10.1002/acr.20632)

Mizner, R. L., Petterson, S. C., Clements, K. E., Zeni, J. A., Irrgang, J. J., & Snyder-Mackler, L. (2011). Measuring functional improvement after total

knee arthroplasty requires both performance-based and patient-report assessments: a longitudinal analysis of outcomes. *The Journal of*

Arthroplasty, 26(5), 728-737.