

References:

1. Hamilton BE, Martin JA, Osterman MHS, Curtin SC, Mathews TJ, Statistics DoV. Births: Final Data for 2014. *Natl Vital Stat Rep* 2015;64.
2. Fanaroff A, Hack M, Walsh M. The NICHD neonatal research network: changes in practice and outcomes during the first 15 years. *Seminars in Neonatology* 2003;27:281-7.
3. Stoll BJ, Hansen NI, Bell EF, et al. Trends in Care Practices, Morbidity, and Mortality of Extremely Preterm Neonates, 1993-2012. *JAMA* 2015;314:1039-51.
4. Stoll BJ, Hansen NI, Bell EF, et al. Neonatal outcomes of extremely preterm infants from the NICHD neonatal research network. *Pediatrics* 2010;126:443-56.
5. Yee WH, Soraisham AS, Shah VS, et al. Incidence and timing of presentation of necrotizing enterocolitis in preterm infants. *Pediatrics* 2012;129:e298-e304.
6. Blakely M, Lally K, McDonald S, Brown R, Barnhart D, Ricketts R. Postoperative outcomes of extremely low birth weight infants with necrotizing enterocolitis or isolated intestinal perforation. *Anal of Surgery* 2005;241:984-94.
7. Day RL, Caliguiri L, Kamenski C, Ehrlich F. Body temperature and survival of premature infants. *Pediatrics* 1964;34:171-81.
8. Hazan J, Maag U, Chesseix P. Association between hypothermia and mortality rate of premature infants - revisited. *Am J Obstet Gynecol* 1991;164:111-2.
9. Buetow KC, Klein SW. Effect of maintenance of "normal" skin temperature of premature infants. *Pediatrics* 1964;34:163-70.
10. Vohra S, Frent G, Campbell V, Abbott M, Whyte R. Effect of polyethylene occlusive skin wrapping on heat loss in very low birth weight infants at delivery: A randomized trial. *The Journal of pediatrics* 1999;134:547-51.
11. Bauer J, Hentschel R, Zahradnik H, Karck U, Linderkamp O. Vaginal delivery and neonatal outcome in extremely-low-birth-weight infants below 26 weeks of gestational age. *Am J Perinatol* 2003;20:181-8.
12. Mank A, van Zanten HA, Meyer MP, Pauws S, Lopriore E, Te Pas AB. Hypothermia in Preterm Infants in the First Hours after Birth: Occurrence, Course and Risk Factors. *PLoS one* 2016;11:e0164817.
13. Yu V, Bajuk J, Astbury J. Perinatal risk factors for necrotizing enterocolitis. *Archives of Disease in Childhood* 1984;59:430-4.
14. Robel-Tillig E, Knupfer M, Pulzer F, Vogtmann C. Blood flow parameters of the superior mesenteric artery as an early predictor of intestinal dysmotility in preterm infants. *Pediatr Radiol* 2004;34:958-62.
15. Knobel R. Fetal and neonatal physiology. *Newborn and Infant Nursing Reviews* 2014;14:45-9.
16. Simbrunner G. Thermoregulation of sick and low birth weight neonates. Germany: Springer-Verlag; 1995.

17. Knobel R, Holditch-Davis D, Schwartz T, Wimmer JE. Extremely low birth weight preterm infants lack vasomotor response in relationship to cold body temperatures at birth. *Journal of Perinatology* 2009;29:814-21.
18. Lyon AJ, Pikaar ME, Badger P, McIntosh N. Temperature control in very low birthweight infants during first five days of life. *Archives in Diseases of Childhood: Fetal/Neonatal Edition* 1997;76:F47-50.
19. Tong VT, Dietz PM, Morrow B, et al. Trends in smoking before, during, and after pregnancy--Pregnancy Risk Assessment Monitoring System, United States, 40 sites, 2000-2010. *MMWR Surveill Summ* 2013;62:1-19.
20. Jeppesen DL, Nielsen SD, Ersboll AK, Valerius NH. Maternal smoking during pregnancy increases the risk of postnatal infections in preterm neonates. *Neonatology* 2008;94:75-8.