

TEXAS WOMAN'S UNIVERSITY

DENTON • DALLAS • HOUSTON

www.twu.edu

CONNECTION OF PARTNER VIOLENCE AND WOMEN'S LOSS OF FERTILITY CONTROL AND IMPACT ON HEALTH

Fuqin Liu, PhD, RN; Judith M. McFarlane, DrPH, RN, FAAN; Sandra Cesario, PhD, MS, BSN, RNC, FAAN

BACKGROUND

- Abused women experience a wide range of adverse health consequences, including poor mental health and impaired reproductive health.
- Women's **lack of fertility control** in an abusive relationship is common.
- Although the association between intimate partner abuse (IPV) and women's poor mental health functioning is well documented, it remains unclear whether women's mental health functioning is linked with compromised fertility control.
 Women's fertility control affects pregnancy outcomes, especially in an abusive relationship.
 Existing studies either focus mainly on the associations between fertility control and unintended pregnancy or examine fertility control as a mediating rather than an independent factor.
 Fertility control has not been examined in relationship to other adverse pregnancy outcomes, such as premature birth.

Figure 1. Fertility Questionnaire

	e trying to learn more about how women make decisions about when to have children, how many children to have, and use of birth I. Thanks for helping us.	NO	YES	66
F1i	Thinking about your relationship with (name of abuser), did you DISCUSS the NUMBER of CHIL- DREN you wanted to have?	0	1	66
F2i	If yes to DISCUSS number of children wanted, did you AGREE on the number of children?	0	1	66
F3i	If yes to AGREE on number of children, did you AGREE on time between pregnancies?	0	1	66
F4i	When you were with () did you DISCUSS using birth control (BC)?	0	1	66
F5i	IF yes to DISCUSS, did you AGREE on type of birth control (condoms, pills, other)?	0	1	66
F6i	When you were with (), was birth control ever used?	0	1	66
F7i	If yes to birth control, who made the decision on type? 1=Abuser; 2=Woman; 3=Joint	Code= 6		66
F8i	Did you ever get pregnant because () would not use BC, such as condoms?	0	1	66
F9i	Did you ever get pregnant because () would not allow you to use birth control?	0	1	66
F10i	Were you ever afraid to talk to () about using birth control?	0	1	66
F11i	Were you ever abused when you talked to () about birth control?	0	1	66
F12i	Were you ever abused when you used birth control?	0	1	66
			•	
Thinkin	ng about the PREGNANCIES you had from (name of abuser)			-
P1i	How many PREGNANCIES did you have with?	1		66
P2i	How many of the pregnancies with were planned?	1		66

OutcomesReproductiveHealth Issues	Women's functioning & safety	Pregnancy outcomes
	Poor mental health Women's Safety	Unintended pregnancy Miscarriage
Fertility control	Study 1: Relationships b/t fertility control, severity of abuse, & poor mental health	Study 2: Links between fertility control and effects on poor pregnancy outcomes

planned pregnancies, who made the plan: 1=Abuser; 2=Woman; 3=Join Code= 66 How many of the pregnancies wit born too soon) baby ecause of abuse? P7i nk the baby died because of the abuse P10i 66 niscarriage, do you think you lost the pregnancy because of abuse? P11i 66 P12i f yes to abortion(s), who made the decision to have the abortion? 1=Abuser; 2=Woman; 3=Joir Code=



RESULTS

Study I:

- Women who reported a pregnancy related to the abuser not using birth control were <u>28</u> times more likely to have had a miscarriage related to abuse (*Odds Ratio* = 28.70, *p* < .05).
- Being abused for using birth control was associated with over <u>8</u> times the risk of having a premature birth (Odds Ratio = 8.34, p < .05).

• Study II:

ANOVA test results with p < .05:

PURPOSE

Study I:

- Can an abused woman's risk for miscarriage be predicted
 - from fertility control items, when controlling for significant covariate?
- Can an abused woman's risk for premature birth be predicted from fertility control items, when controlling for significant covariate?

Study II:

• The purpose of the study is to examine the relationships among fertility control, severity of abuse, and poor mental health among abused women.

METHODS

- Parent study: A longitudinal study to examine the long-term health and functioning consequences of IPV among 150 first-time users of safe shelters and 150 first-time users of the justice system following an abusive relationship.
- Repeated measures every 4 months.
- Measures assessed severity of abuse; danger for murder; mental health symptoms of depression, anxiety, somatization, and PTSD.
- At 32 months, 282 women completed the investigatordeveloped fertility control questionnaire.
- Statistical procedures: ANNOVA & logistic regression.

Fertility Control Items	Severity of Violence	Poor Mental Health
1.Discuss # of kids		1. PTSD
2. Agree on number of children?	2. Threats	
3. Agree on time b/t pregnancies?		3. PTSD
4. Used birth control?	4. Danger	4. Depression & PTSD
5. Who decided on birth control?		5. Depression & PTSD
5-1. Abuser		
5-2. Woman		6. PTSD
5-3. Joint		
6.Afraid to talk to abuser about birth control?		
7. Abused because used birth control?	7. Physical Abuse	

DISCUSSION

- Compromised fertility control is **associated with** adverse pregnancy outcomes, such as preterm birth and miscarriage.
- Controlling experiences around one's reproductive health is **associated with** what appears to be more severe forms of abuse (i.e. increased lethality risk). Lack of control over fertility is **associated with** poorer mental health.

ACKNOWLEDGMENTS

Funded by Houston Foundation

